

## INSTRUCTIONS

This question paper contains total 150 questions divided into four parts:

Part I : Physics Q. No. 1 to 40

Part II : Chemistry Q. No. 41 to 80

Part III : (A) English Proficiency Q. No. 81 to 95





(B) Logical Reasoning Q. No. 96 to 105

Part IV : Mathematics Q. No. 106 to 150


All questions are multiple choice questions with four options, only one of them is correct. Each correct answer awarded 3 marks and -1 for each incorrect answer.

### PART - I : PHYSICS

1. Which one of the following graphs represents the variation of electric field with distance  $r$  from the centre of a charged spherical conductor of radius  $R$ ?



Duration of paper-3 Hours



(c)  $3\Omega$

(d)  $4\Omega$

2. If  $\vec{E}$  and  $\vec{B}$  are the electric and magnetic field vectors of e.m. waves then the direction of propagation of e.m. wave is along the direction of  $\vec{E} \times \vec{B}$ .

3. The young's modulus of a wire of length  $L$  and radius  $r$  is  $Y$  N/m<sup>2</sup>. If the length and radius are

A

(a)  $1W$

(b)  $2W$

(a)  $E \cdot B$

(b)  $B$

(c)  $E \cdot B$

(d) None of these

reduced to  $L/2$  and  $r/2$ , then its young's modulus will be

- (a)  $Y/2$  (b)  $Y$  (c)  $2Y$  (d)  $4Y$

4. Twelve resistors each of resistance  $16 \Omega$  are connected in the circuit as shown. The net resistance between A and B is

5. The time period of a satellite of earth is 5 hours. If the separation between the earth and the satellite is increased to 4 times the previous value, the new time period will become

- (a) 10 hours (b) 80 hours  
(c) 40 hours (d) 20 hours

6. Two trains are moving towards each other with speeds of  $20 \text{ m/s}$  and  $15 \text{ m/s}$  relative to the ground. The first train sounds a whistle of frequency  $600 \text{ Hz}$ . The frequency of the whistle heard by a passenger in the second train before the train meets, is (the speed of sound in air is

$340 \text{ m/s}$ )


- (a)  $600 \text{ Hz}$  (b)  $585 \text{ Hz}$   
(c)  $645 \text{ Hz}$  (d)  $666 \text{ Hz}$

7. You are asked to design a shaving mirror assuming that a person keeps it  $10 \text{ cm}$  from his face and views the magnified image of the face at the closest comfortable distance of  $25 \text{ cm}$ . The radius of curvature of the mirror would then be :

- (a)  $60 \text{ cm}$  (b)  $-24 \text{ cm}$   
(c)  $-60 \text{ cm}$  (d)  $24 \text{ cm}$

8. A block is kept on a frictionless inclined surface with angle of inclination ' $\alpha$ '. The incline is given an acceleration ' $a$ ' to keep the block stationary. Then ' $a$ ' is equal to

- (a)  $g \operatorname{cosec} \alpha$   
(b)  $g/\tan \alpha$   
(c)  $g \tan \alpha$   
(d)  $g$



9. With the increase in temperature, the angle of contact

- (a) decreases  
(b) increases  
(c) remains constant  
(d) sometimes increases and sometimes decreases

10. Forward biasing is that in which applied voltage

- (a) increases potential barrier  
(b) cancels the potential barrier  
(c) is equal to  $1.5 \text{ volt}$   
(d) None of these

11. Number of significant figures in expression  $4.327g$

$2.51 \text{ cm}^3$  is

- (a) 2 (b) 4 (c) 3 (d) 5

12. The ratio of the specific heats  $\frac{C_p}{C_v} = \gamma$  in terms of degrees of freedom ( $n$ ) is given by

(a)  $\frac{n+2}{n}$  (b)  $\frac{n}{n+2}$

(c)  $\frac{n+2}{n-2}$  (d)  $\frac{n-2}{n+2}$

13. A stone is thrown with a velocity  $u$  making an angle  $\theta$  with the horizontal. The horizontal distance covered by its fall to ground is maximum

when the angle  $\theta$  is equal to

- (a)  $0^\circ$  (b)  $30^\circ$  (c)  $45^\circ$  (d)  $90^\circ$

14. A ball of mass  $150 \text{ g}$ , moving with an acceleration  $20 \text{ m/s}^2$ , is hit by a force, which acts on it for  $0.1 \text{ sec}$ . The impulsive force is

- (a)  $0.5 \text{ N}$  (b)  $0.1 \text{ N}$  (c)  $0.3 \text{ N}$  (d)  $1.2 \text{ N}$

15. A man drags a block through  $10 \text{ m}$  on rough surface ( $\mu = 0.5$ ). A force of  $\sqrt{5} \text{ kN}$  acting at  $30^\circ$  to the horizontal. The work done by applied force is

- (a) zero (b)  $7.5 \text{ kJ}$  (c)  $5 \text{ kJ}$  (d)  $10 \text{ kJ}$

16. A force of  $(2\hat{i} + 3\hat{j} + 4\hat{k}) \text{ N}$  acts on a body for 4 second, produces a displacement of

$(3\hat{i} + 4\hat{j} + 5\hat{k}) \text{ m}$ . The power used is

- (a)  $9.5 \text{ W}$  (b)  $7.5 \text{ W}$  (c)  $6.5 \text{ W}$  (d)  $4.5 \text{ W}$

17. The Earth is assumed to be a sphere of radius  $R$ . A platform is arranged at a height  $R$  from the surface of the Earth. The escape velocity of a body from this platform is  $f v$ , where  $v$  is its escape velocity from the surface of the Earth. The value of  $f$  is

(a)  $\frac{1}{3}$  (b)  $\frac{1}{2}$  (c)  $2$  (d)  $\frac{1}{\sqrt{2}}$

18. Kepler's second law regarding constancy of areal velocity of a planet is a consequence of the law of conservation of


- (a) Energy
- (b) Angular momentum
- (c) Linear momentum
- (d) None of these

19. Water is flowing through a horizontal tube having cross-sectional areas of its two ends being  $A$  and  $A'$  such that the ratio  $A/A'$  is 5. If the pressure difference of water between the two ends is  $3 \times 10^5 \text{ N m}^{-2}$ , the velocity of water with which it enters the tube will be (neglect gravity effects)

- (a)  $5 \text{ m s}^{-1}$  (b)  $10 \text{ m s}^{-1}$
- (c)  $25 \text{ m s}^{-1}$  (d)  $50\sqrt{10} \text{ m s}^{-1}$

20. A thermodynamic system is taken from state A to B along  $ACB$  and is brought back to A along  $BDA$  as shown in the  $PV$  diagram. The net work done during the complete cycle is given by the area

- (a)  $P_1ACBP_2P_1$
- (b)  $ACBB'A'A$
- (c)  $ACBDA$
- (d)  $ADBB'A'A$



21. A boat crosses a river from port A to port B, which are just on the opposite side. The speed of the water is  $V_w$  and that of boat is  $V_B$  relative to still water. Assume  $V_w = 2V_B$ . What is the time taken by the boat, if it has to cross the river directly on the AB line [ $D$  = width of the river]

2D

3D

(a)  $\frac{\sqrt{D}}{V_B \sqrt{2}}$  (b)  $\frac{\sqrt{D}}{2V_B}$

(c)  $\frac{D}{V_B \sqrt{2}}$  (d)  $V_B$

22. Two springs, of force constants  $k_1$  and  $k_2$  are connected to a mass  $m$  as shown. The frequency of oscillation of the mass is  $f$ . If both  $k_1$  and  $k_2$  are made four times their original values, the frequency of oscillation becomes  $k$



- (a)  $2f$  (b)  $f/2$  (c)  $f/4$  (d)  $4f$

23. When a potential difference  $V$  is applied across a conductor at a temperature  $T$ , the drift velocity of electrons is proportional to

- (a)  $\sqrt{V}$  (b)  $V$  (c)  $\sqrt{T}$  (d)  $T$

24. The amplitude of a damped oscillator becomes  $\frac{1}{e}$  in 2 seconds. If its amplitude after  $n$


seconds is  $\frac{1}{e^n}$  times the original amplitude, the value of  $n$  is

- (a)  $3^2$  (b)  $3^3$  (c)  $\sqrt[3]{3}$  (d)  $2^3$

25. The angular speed of the electron in the  $n^{\text{th}}$  orbit of Bohr hydrogen atom is (a) directly proportional to  $n$

- (b) inversely proportional to  $n$
- (c) inversely proportional to  $n^2$
- (d) inversely proportional to  $n^3$

26. In the given figure, the charge on  $3 \mu\text{F}$  capacitor is



- (a)  $10 \mu\text{C}$  (b)  $15 \mu\text{C}$
- (c)  $30 \mu\text{C}$  (d)  $5 \mu\text{C}$

27. Two bodies A and B are placed in an evacuated vessel maintained at a temperature of  $27^\circ\text{C}$ . The temperature of A is  $327^\circ\text{C}$  and that of B is


227°C. The ratio of heat loss from A and B is about

- (a) 2 : 1 (b) 1 : 2 (c) 4 : 1 (d) 1 : 4
28. If a rigid body is rotating about an axis with a constant velocity, then
- (a) Velocity, Angular velocity of all particles will be same  
 (b) Velocity, Angular velocity of all particles will be different  
 (c) Velocity of all particles will be different but angular velocity will be same.  
 (d) Angular velocity of all particles will be different but velocity will be same.
29. The fundamental frequency of an open organ pipe is 300 Hz. The first overtone of this pipe has same frequency as first overtone of a closed organ pipe. If speed of sound is 330 m/s, then the length of closed organ pipe is
- (a) 41 cm (b) 30 cm (c) 45 cm (d) 35 cm
30. In Young's experiment, the distance between the slits is reduced to half and the distance between the slit and screen is doubled, then the fringe width (a) will not change  
 (b) will become half  
 (c) will be doubled  
 (d) will become four times
31. If a rolling body's angular momentum changes by 20 SI units in 3 seconds, by a constant torque. Then find the torque on the body
- (a) 20/3 SI units (b) 100/3 SI units  
 (c) 20 SI units (d) 5 SI units
32. Charge Q is distributed to two different metallic spheres having radii  $x$  and  $2x$  such that both spheres have equal surface charge density, then charge on large sphere is
- (a)  $\frac{4Q}{5}$  (b)  $\frac{Q}{5}$  (c)  $\frac{3Q}{5}$  (d)  $\frac{5Q}{4}$
33. In an LR circuit  $f = 50$  Hz,  $L = 2$  H,  $E = 5$  volts,  $R = 1$  W then energy stored in inductor is
- (a) 50 J (b) 25 J  
 (c) 100 J (d) None of these
34. A straight wire of length 0.5 metre and carrying a current of 1.2 ampere is placed in uniform magnetic field of induction 2 tesla. The


magnetic field is perpendicular to the length of the wire.

The force on the wire is

- (a) 2.4 N (b) 1.2 N (c) 3.0 N (d) 2.0 N
35. A man drives a car from station B towards station A at speed 60 km/h. A car leaves station A at speed 60 km/h every 10 min. The distance between A and B is 60 km. The car travels at the speed of 60 km/h. A man drives a car from B towards A at speed of 60 km/h. If he starts at the moment when first car leaves the station B, then how many cars would be met on the route ?
- (a) 4 (b) 7 (c) 9 (d) 12
36. In rotatory motion, linear velocities of all the particles of the body are
- (a) same (b) different  
 (c) zero (d) cannot say
37. If  $x$ ,  $v$  and  $a$  denote the displacement, the velocity and the acceleration of a particle executing simple harmonic motion of time period  $T$ , then, which of the following does not change with time?
- (a)  $aT/x$  (b)  $aT + 2pv$   
 (c)  $aT/v$  (d)  $a^2T^2 + 4p^2v^2$
38. A conducting wire frame is placed in a magnetic field which is directed into the paper. The magnetic field is increasing at a constant rate. The directions of induced current in wires AB and CD are






- (a) B to A and D to C (b) A to B and C to D  
 (c) A to B and D to C (d) B to A and C to D
39. Find the acceleration of block A and B. Assume pulley is massless.



$$(a) \frac{10}{13} \text{ g}, \frac{5}{13} \text{ g} \quad (b) \frac{1}{13} \text{ g}, \frac{5}{13} \text{ g}$$





$$(c) \frac{9}{13} \text{ g}, \frac{11}{13} \text{ g} \quad (d) \frac{13}{10} \text{ g}, \frac{13}{5} \text{ g}$$

40. The nuclei of which one of the following pairs of nuclei are isotones?



## PART - II : CHEMISTRY

41. Plots showing the variation of the rate constant ( $k$ ) with temperature ( $T$ ) are given below. The plot that follows Arrhenius equation is



42. 3.6 g of oxygen is adsorbed on 1.2 g of metal powder. What volume of oxygen adsorbed per gram of the adsorbent at 1 atm and 273 K?

(a)  $0.19 \text{ L g}^{-1}$

(b)  $1 \text{ L g}^{-1}$

(c) 2.1 L  $\text{g}^{-1}$  (d) None of these

43. In the purification of impure nickel by Mond's process, metal is purified by :

(a) Electrolytic reduction

(b) Vapour phase thermal decomposition

(c) Thermite reduction

(d) Carbon reduction

44. When chlorine water is added to an aqueous solution of sodium iodide in the presence of chloroform, a violet colouration is obtained. On adding more of chlorine water and vigorous shaking, the violet colour disappears. This shows the conversion of ..... into .....



45. In the clathrates of xenon with water, the nature of bonding between xenon and water molecule is





(a) covalent

(b) hydrogen bonding


(c) coordinate

(d) dipole-induced dipole

46. The electronic configurations of Eu (Atomic No. 63), Gd (Atomic No. 64) and Tb (Atomic No. 65) are




47. Which of the following carbonyls will have the strongest C – O bond ?




48. How many chiral compounds are possible on monochlorination of 2-methyl butane ?






(a) 8 (b) 2 (c) 4 (d) 6

49. Which of the following are intermediates in the reaction of excess of  $\text{CH}_3\text{MgBr}$  with









÷









170°C

What is Z?



51. Which of the following is the strongest base ?



52. Which of the following does not reduce Benedict's solution?

- (a) Glucose (b) Fructose  
 (c) Sucrose (d) Aldehyde

53. General formula of solid in zinc blende structure is:

- (a)  $\text{AB}_2$  (b)  $\text{AB}_3$  (c)  $\text{AB}$  (d)  $\text{A}_2\text{B}$

54. Glycine in alkaline solution exists as \_\_\_\_\_ and migrates to \_\_\_\_\_. (a) Cation, cathode

- (b) Neutral, anode  
 (c) Zwitter ion, cathode  
 (d) anion, anode

55. Product on reaction of ethanamide with phosphorus pentoxide is:

- (a) ethanamine  
 (b) acetonitrile  
 (c) ethanol  
 (d) ethane isonitrile

56.  $K_a$  of  $\text{HX}$  is  $10^{-5}$ , then find concentration of  $\text{H}_3\text{O}^+$  when equal volumes of 0.25M  $\text{HX}$  and 0.05 M  $\text{NaOH}$  are mixed.

- (a)  $4 \times 10^{-5} \text{ M}$  (b)  $6 \times 10^{-5} \text{ M}$   
 (c)  $8 \times 10^{-3} \text{ M}$  (d)  $2 \times 10^{-5} \text{ M}$

57. Net cell reaction of  $\text{Pt} | \text{H}_2 (640 \text{ mm}) | \text{HCl} | \text{H}_2 (510 \text{ mm}) | \text{Pt}$ .

- (a) 0.89 V (b) 0.93 V  
 (c)  $2.91 \times 10^{-3} \text{ V}$  (d)  $2.5 \times 10^{-2} \text{ V}$

58. Which of the following has zero net dipole moment?

- (a)  $\text{XeF}_4$  (b)  $\text{BrF}_3$  (c)  $\text{ClF}_3$  (d)  $\text{SF}_4$

59. Which of the following element has the highest ionisation enthalpy?

- (a) Boron (b) Aluminium  
 (c) Germanium (d) Thallium

60. Out of the elements with atomic number 7, 8, 9, 13 which has the smallest size and highest ionization enthalpy?

(a) 7 (b) 8 (c) 9 (d) 13

61. Which one is classified as a condensation polymer?

(a) Dacron (b) Neoprene  
(c) Teflon (d) Acrylonitrile


62. Which of the following compounds is not an antacid?

(a) Phenelzine (b) Ranitidine  
(c) Aluminium hydroxide (d) Cimetidine

63. Mole fraction of the solute in a 1.00 molal aqueous solution is

(a) 0.1770 (b) 0.0177 (c) 0.0344 (d) 1.7700

64. The IUPAC name of the following compound is






(a) *trans*-2-chloro-3-iodo-2-pentene (b) *cis*-3-iodo-4-chloro-3-pentene

(c) *trans*-3-iodo-4-chloro-3-pentene


(d) *cis*-2-chloro-3-iodo-2-pentene

65. Most stable carbocation among the following is:



(d)  $\text{CH}_3^+$

66. Which is correct for the following changes?



Me

(a) X is Lindlar Catalyst, B is *cis*-2-butene

(b) A is 2-butyne, X is Na-liq.  $\text{NH}_3$

(c) B is *trans*-2-butene, X is Na-liq.  $\text{NH}_3$

(d) A is 2-butene, X is  $\text{SeO}_2$

67. The stability of +1 oxidation state among Al, Ga, In and Tl increases in the sequence:

(a)  $\text{Ga} < \text{In} < \text{Al} < \text{Tl}$  (b)  $\text{Al} < \text{Ga} < \text{In} < \text{Tl}$

(c)  $\text{Tl} < \text{In} < \text{Ga} < \text{Al}$

(d)  $\text{In} < \text{Tl} < \text{Ga} < \text{Al}$

68. Which of the following alkaline earth metal hydroxides is amphoteric in character?

(a)  $\text{Be}(\text{OH})_2$  (b)  $\text{Ca}(\text{OH})_2$

(c)  $\text{Sr}(\text{OH})_2$  (d)  $\text{Ba}(\text{OH})_2$

69. Which reaction shows oxidising nature of  $\text{H}_2\text{O}_2$ ?

(a)  $\text{H}_2\text{O}_2 + 2\text{KI} \rightarrow 2\text{KOH} + \text{I}_2$

(b)  $\text{Cl}_2 + \text{H}_2\text{O}_2 \rightarrow 2\text{HCl} + \text{O}_2$

(c)  $\text{H}_2\text{O} + \text{Ag}_2\text{O} \rightarrow 2\text{Ag} + \text{H}_2\text{O} + \text{O}_2$

(d)  $\text{NaClO} + \text{H}_2\text{O}_2 \rightarrow \text{NaCl} + \text{H}_2\text{O} + \text{O}_2$

70.  $a\text{K}_2\text{Cr}_2\text{O}_7 + b\text{KCl} + c\text{H}_2\text{SO}_4 \rightarrow 4\text{K}_2\text{SO}_4 + x\text{Cr}_2\text{O}_3$

$\text{Cl}_2 + y\text{KHSO}_4 + z\text{H}_2\text{O}$

The above equation balances when




(a)  $a = 2, b = 4, c = 6$  and  $x = 2, y = 6, z = 3$

(b)  $a = 4, b = 2, c = 6$  and  $x = 6, y = 2, z = 3$

(c)  $a = 6, b = 4, c = 2$  and  $x = 6, y = 3, z = 2$

(d)  $a = 1, b = 4, c = 6$  and  $x = 2, y = 6, z = 3$

71. For the reactions



$K_c$  for the reaction  $\text{A} \rightleftharpoons \text{D}$  is

(a)  $2 \times 4 \times 6$  (b)  $\frac{2 \times 4}{6}$   
(c)  $2 + 4 + 6$  (d)  $\frac{4 \times 6}{2}$

72. Which of the following will always lead to a non-spontaneous change? (a) DH and DS both +ve  
 (b) DH is -ve DS both +ve (c) DH and DS both -ve  
 (d) DH is +ve DS both -ve
73. The densities of two gasses are in the ratio of 1: 16. The ratio of their rates of diffusion is  
 (a) 16:1 (b) 4:1 (c) 1:4 (d) 1:16
74. In the reaction  $2\text{PCl}_5 \rightleftharpoons \text{PCl}_4^+ + \text{PCl}_6^-$ , the change in hybridisation is from  
 (a)  $sp^3d$  to  $sp^3$  and  $sp^3d^2$   
 (b)  $sp^3d$  to  $sp^2$  and  $sp^3$   
 (c)  $sp^3d$  to  $sp^3d^2$  and  $sp^3d^3$   
 (d)  $sp^3d^2$  to  $sp^3$  and  $sp^3d$
75. The group having isoelectronic species is:  
 (a)  $\text{O}^{2-}$ ,  $\text{F}^-$ ,  $\text{Na}^+$ ,  $\text{Mg}^{2+}$  (b)  $\text{O}^-$ ,  $\text{F}^-$ ,  $\text{Na}$ ,  $\text{Mg}^+$   
 (c)  $\text{O}^{2-}$ ,  $\text{F}^-$ ,  $\text{Na}$ ,  $\text{Mg}^{2+}$   
 (d)  $\text{O}^-$ ,  $\text{F}^-$ ,  $\text{Na}^+$ ,  $\text{Mg}^{2+}$
76. 100 mL  $\text{O}_2$  and  $\text{H}_2$  kept at same temperature and pressure. What is true about their number of molecules (a)  $\text{NO}_2 > \text{NH}_2$   
 (b)  $\text{NO}_2 < \text{NH}_2$   
 (c)  $\text{NO}_2 = \text{NH}_2$   
 (d)  $\text{NO}_2 + \text{NH}_2 = 1$  mole
77. If  $m_A$  gram of a metal A displaces  $m_B$  gram of another metal B from its salt solution and if the equivalent mass are  $E_A$  and  $E_B$  respectively then equivalent mass of A can be expressed as:  
 $m^{\frac{A}{E_B}}$  (a)  $E_A = m_B$   
 (b)  $E_A = m_A \cdot m_B$   
 (c)  $E_A = m_A \cdot \frac{E_B}{m_B}$   
 (d)  $E_A = m_B \sqrt{m^{\frac{A}{E_B}} E_B}$

78. Which one of the following set of quantum numbers is not possible for 4p electron?





(a)  $n = 4, l = 1, m = -1, m_s = +\frac{1}{2}$

(b)  $n = 4, l = 1, m = 0, m_s = +\frac{1}{2}$

(c)  $n = 4, l = 1, m = 2, m_s = +\frac{1}{2}$

(d)  $n = 4, l = 1, m = -1, m_s = -\frac{1}{2}$

79. Which of the following radial distribution graphs correspond to  $l = 2$  for the H atom ?



80. Which of the following is paramagnetic?

- (a)  $\text{B}_2$  (b)  $\text{C}_2$  (c)  $\text{N}_2$  (d)  $\text{F}_2$

**PART - III (A): ENGLISH PROFICIENCY**

**DIRECTIONS (Qs. 81-83) :** In the following questions below, out of the four alternatives, choose



*the one which best expresses the meaning of the given word.*

- 
- 81.** Garrulous  
(a) Talkative (b) Sedative  
(c) Cocative (d) Positive
- 82.** Tinsel  
(a) Tinkle (b) Decoration  
(c) Tin (d) Colourful
- 83.** Labyrinth  
(a) Meandering (b) Rotating  
(c) Pacing (d) Wriggling

---

**DIRECTIONS (Qs. 84-86) :** *In the following questions, choose the word opposite in meaning to the given word.*

- 
- 84.** Knack :  
(a) Talent (b) Dullness  
(c) Dexterity (d) Balance
- 85.** Pernicious :  
(a) Prolonged (b) Ruinous  
(c) Ruthless (d) Beneficial
- 86.** Opulence :  
(a) Luxury (b)  
Transparency  
(c) Weath (d) Poverty

---

**DIRECTIONS (Qs. 87-90) :** *Read the passage carefully and choose the best answer to each question out of the four alternatives and mark it by blackening the appropriate circle [·].*

---

Like watering a plant, we grow our friendships [and all our relationships) by running them. Friendships need the same attention as other relationships. If they are to continue. These relationships can be delightfully non-judgemental, supportive, understanding and fun.

Sometimes a friendship can bring out the positive side that you never show in any other relationship. This may be because the pressure of playing a 'role' (daughter, partner or child) is removed. With a friend you are to be yourself and free to change. Of course, you are free to do this in all other relationships as well, but in friendships you get to have lats of rehearsals and discussion about changes as you experience them. It is an unconditional experience where you receive as much as you give. You can explain yourself to a friend openly without the fear of hurting a family member. How do friendships grow ? The answer is simple. By revealing yourself; being attentive: remembering what is most showing empathy; seeing the world through the eyes of your friend, you will understand the value of friendship. All this means learning to accept a person from a completely different family to your own or perhaps someone from a completely different cultural background. This is the way we learn tolerance. In turn we gain tolerance and acceptance for our own differences.

- 87.** In good friendships, we (a) give and receive.  
(b) neither give nor receive.  
(c) only give.  
(d) only receive.
- 88.** Empathy means  
(a) someone else's misfortunes  
(b) the ability to share and understand another feelings.  
(c) skill and efficiency  
(d) ability to do something
- 89.** Through strong friendships, we gain (a) only acceptance.  
(b)only attention.  
(c)acceptance and tolerance. (d)  
only tolerance.
- 90.** Friendships and relationships grow when they are  
(a) compared (b) divided  
(c) favoured (d) nurtured
-

**DIRECTIONS (Qs. 91-92) :** In the following questions, sentences are given with blanks to be filled with an appropriate word(s). Four alternatives are suggested for each question. Choose the correct alternative out of the four as your answer.

91. There are not solitary, free-living creatures ; every form of life is \_\_\_\_\_ other forms.  
 (a) dependent on (b) parallel to  
 (c) overshadowed by (d) segregated from
92. I'll take \_\_\_\_\_ now as I have another's appointment some where else.  
 (a) departure (b) your leave  
 (c) permission (d) leave from work


**DIRECTIONS (Qs. 93-95) :** In the following questions, some parts of the sentences have errors and some are correct. Find out which part of a sentence has an error. The number of that part is the answer. If a sentence is free from error, then your answer is (d). i.e., No error.

93. When one hears of the incident (a)/about the plane crash (b)/ he feels very sorry. (c)/ No error (d)
94. I went there (a)/ with a view to survey (b)/ the entire procedure. (c)/ No error (d)
95. It had laid (a)/ in the closet (b)/ for a week before we found it. (c)/ No error (d)


**PART - III (B) : LOGICAL REASONING**

**DIRECTIONS (Qs. 96 & 97) :** In the following questions, which answer figure will complete the question figure?


96. Question Figures :




Answer figures :



97. Question Figure:




Answer Figure:




98. A piece of paper is folded and cut/punched as shown below in the question figures. From the given answer figures, indicate how it will appear when opened.

Question figures:



Answer figures:



99. Select the related word from the given alternatives:  
 Medicine : Patient :: Education : ?  
 (a) Teacher (b) School  
 (c) Student (d) Tuition
100. Choose the correct alternative from the given ones that will complete the series.  
 A3E, F5J, K7O, \_\_\_\_\_  
 (a) Q11T (b) Q9V  
 (c) P9T (d) P11T
101. Which one of the following numbers lacks the common property in the series?

81, 36, 25, 9, 5, 16

(a) 5 (b) 9

(c) 36 (d) 25

102. In a certain code language, "TIRED" is written as "56" and "BRAIN" is written as "44". How is "LAZY" written in that code language?

(a) 64 (b) 61

(c) 58 (d) 43

103. Select the missing number from the given response.

8	7	6
8	7	6
88	77	?
5632	3773	3132

(a) 66 (b) 87 (c) 78 (d) 76

104. Which one of the following diagrams best depicts the relationship among Human Society - Youth Club, Political Party and Youths ?



(a)(b)



(c)(d)



105. Among her children, Ganga's favourites are Ram and Rekha. Rekha is the mother of Sharat, who is loved most by his uncle Mithun. The head of the family is Ram Lal, who is succeeded by his sons Gopal and Mohan. Gopal and Ganga have been married for 35 years and have 3 children.

What is the relation between Mithun and Mohan?

(a) Uncle (b) Son

(c) Brother (d) No relation

## PART - IV : MATHEMATICS

106. If  $x \cos a + y \sin a = P$  is a tangent to the ellipse

$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , then

$\frac{P}{a} = \frac{P}{b}$ , then

$a$

(a)  $a \cos a + b \sin a = P^2$

(b)  $a \sin a + b \cos a = P^2$

(c)  $a^2 \cos^2 a + b^2 \sin^2 a = P^2$

(d)  $a^2 \sin^2 a + b^2 \cos^2 a = P^2$

107. If  $a_1, a_2, a_3, \dots, a_n$  are in A.P. where  $a_i > 0$  for all  $i$ , then

$$\frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} =$$

(a)  $\frac{n+1}{\sqrt{a_1} + \sqrt{a_n}}$  (b)  $\frac{n}{\sqrt{a_1} + \sqrt{a_n}}$

(c)  $\frac{n-1}{\sqrt{a_1} + \sqrt{a_n}}$  (d) none of these

108. In order to solve the differential equation

$\frac{dy}{dx}$

$$x \cos x + y(x \sin x + \cos x) = 1$$

the integrating factor is:

(a)  $x \cos x$  (b)  $x \sec x$

(c)  $x \sin x$  (d)  $x \operatorname{cosec} x$

109. Equation of two straight lines are  $x-1 = y-2 = z-3$  and  $x-4 = y-1 = z$ .

$$x-3 \text{ and } x-4 = y-1 = z.$$

2 3 4 5 2

Then

(a) The lines are non-coplanar

(b) The lines are parallel and distinct

(c) The lines intersect in unique point

(d) The lines are coincident

110. The equation of the curve passing through the

point  $(a, -1)$  and satisfying the

differential equation  $y - x \frac{dy}{dx} = a y^2$

$+ \frac{dy}{dx} = 0$  is

(a)  $(x+a)(1+ay) = -4a y^2$  (b)  $(x+a)(1-ay) = 4a^2 y$

(c)  $(x+a)(1-ay) = -4a^2 y$

(d) None of these

111. The locus of the mid-point of a chord of the circle  $x^2 + y^2 = 4$ , which subtends a right angle at the

origin is

- (a)  $x + y = 2$  (b)  $x^2 + y^2 = 1$   
 (c)  $x^2 + y^2 = 2$  (d)  $x + y = 1$

112. With the usual notation  $\int_1^{\infty} ([x^2] - [x]^2) dx$  is equal to

- (a)  $4 + 2\sqrt{3}$  (b)  $4 - 2\sqrt{3}$   
 (c)  $4 - 2\sqrt{3}$  (d) none of these

113.

$$\frac{1}{1 + \sin A + \cos A} = \frac{1 + \sin A - \cos A}{2}$$

- (a)  $\sin \frac{A}{2}$  (b)  $\cos \frac{A}{2}$   
 (c)  $\tan \frac{A}{2}$  (d)  $\cot \frac{A}{2}$

114.

If  $x \sqrt{y} + y \sqrt{x} = \frac{x+1}{2}$

then  $\frac{dy}{dx} =$

- (a)  $\frac{x}{1+x}$  (b)  $\frac{x}{1+x}$   
 (c)  $\frac{-1}{2}$  (d)  $1$

115.

If  $f(x) = 3x^4 + 4x^3 - 12x^2 + 12$ , then  $f(x)$  is

- (a) increasing in  $(-\infty, -2)$  and in  $(0, 1)$

- (b) increasing in  $(-2, 0)$  and in  $(1, \infty)$   
 (c) decreasing in  $(-2, 0)$  and in  $(0, 1)$   
 (d) decreasing in  $(-\infty, -2)$  and in  $(1, \infty)$

116.

$$\begin{matrix} x & y & x & y \\ - & - & + & + \\ 2 & 4 & 3 & 2 \end{matrix}$$

Then number of possible solutions are :

- (a) Zero (b) Unique  
 (c) Infinite (d) None of these

117.

The distance of a point  $(2, 5, -3)$  from the plane  $r \times (6i - 3j + 2k) = 4$  is

(a)  $\frac{13}{5}$  (b)  $\frac{13}{7}$   
 (c)  $\frac{13}{5}$  (d)  $\frac{37}{7}$


118.

The value of definite integral  $\int_0^p \log(\tan x) dx$  is

(a)  $\frac{p}{4}$  (b)  $\frac{p}{2}$   
 (c)  $\frac{p}{2}$  (d)  $p$

119.

For the following feasible region, the linear constraints are



- (a)  $x \geq 0, y \geq 0, 3x + 2y \leq 12, x + 3y \leq 11$   
 (b)  $x \geq 0, y \geq 0, 3x + 2y \leq 12, x + 3y \leq 11$   
 (c)  $x \geq 0, y \geq 0, 3x + 2y \leq 12, x + 3y \leq 11$

120. (d) None of these  
The general solution of differential equation

$$(e^x + 1) y dy = (y + 1) e^x dx$$

- (a)  $(y + 1) = k(e^x + 1)$   
 (b)  $y + 1 = e^x + 1 + k$   
 (c)  $y = \log \{k(y + 1) (e^x + 1)\}$

(d)  $y = \log \left[ \frac{e^{y^2 + 1} (y + 1) + k}{e^{y^2 + 1} + k} \right]$

121. What is the slope of the normal at the point  $(at^2, 2at)$  of the parabola  $y^2 = 4ax$ ?

- (a)  $-\frac{1}{t}$  (b)  $t$   
 (c)  $-t$  (d)  $-\frac{1}{t}$

122.  $\int_0^{\pi/2} x \sin^2 x \cos^2 x dx$  is equal to

- (a)  $\frac{\pi^2}{32}$  (b)  $\frac{\pi^2}{16}$   
 (c)  $\frac{\pi}{32}$  (d) None of these

123.  $\left| \begin{matrix} 6i - 3i & 1 \\ 1 & 1 \end{matrix} \right|$  If  $4 - 3i + ix = \sqrt{-1}$ , where  $i = -1$ , then

- 203  $i$  what is  $x$  equal to?  
 (a) 3 (b) 2  
 (c) 1 (d) 0

124. The limit  $\lim_{x \rightarrow 0} \frac{\log_e(21+x) + x}{1+x}$

- (a) is equal to  $\frac{1}{2}$  (b) is equal to  $-\frac{1}{2}$   
 (c) is equal to 2 (d) does not exist

125. If  $2 \cos^2 x + 3 \sin x - 3 = 0$ ,  $0 \leq x \leq 180^\circ$ , then  $x =$

- (a)  $30^\circ, 90^\circ, 150^\circ$  (b)  $60^\circ, 120^\circ, 180^\circ$   
 (c)  $0^\circ, 30^\circ, 150^\circ$  (d)  $45^\circ, 90^\circ, 135^\circ$

126. If the number of available constraints is 3 and the number of parameters to be optimized is 4, then (a) The objective function can be optimized

- (b) The constraint are short in number  
 (c) The solution is problem oriented  
 (d) None of these

127.  $\sqrt{\frac{1}{2}}$  If  $y = \tan^{-1} \frac{e^x - e^{-x}}{e^x + e^{-x}}$ , then  $y'(1)$  is equal to

- (a) 0 (b)  $\frac{1}{2}$  (c) -1 (d)  $-\frac{1}{4}$

128. The maximum area of rectangle inscribed in a circle of diameter R is

- (a)  $\frac{R^2}{2}$  (b)  $\frac{R^2}{4}$   
 (c)  $\frac{R^2}{8}$  (d)  $\frac{R^2}{16}$

129. If A and B are two events, such that

$$P(A \cap B) = \frac{3}{8}, P(A \cup B) = \frac{1}{2}, P(A)^c = \frac{2}{3}$$

where  $A^c$  stands for the complementary event of A, then  $P(B)$  is given by:

- (a)  $\frac{1}{3}$  (b)  $\frac{2}{3}$   
 (c)  $\frac{1}{9}$  (d)  $\frac{2}{9}$

130.  $\lim_{x \rightarrow 0} \frac{e^{1/x} - e^{-1/x}}{x^k}$ , If  $f(x) = e^{1/x} + e^{-1/x}$  then  $\lim_{x \rightarrow 0} f(x)$  exists if  
 (a)  $f$  is continuous at  $x$ , when  $k = 0$  (b)  $f$  is not continuous at  $x = 0$  for any real  $k$ . (c)  $\lim_{x \rightarrow 0} f(x)$  exist infinitely  $x \neq 0$   
 (d) None of these

131.  $\int_0^{\pi/2} \cos^{-1}(\sin 2x) dx$  is equal to  $\frac{1}{2} \ln k$   
 (a)  $\frac{1}{2} \ln(x^2 + k)$  (b)  $\frac{1}{2} \ln(x^2 + 1)$   
 (c)  $\frac{1}{2} \ln(x + k)$  (d) None of these

132. The equation of chord of the circle  $x^2 + y^2 = 8x$  bisected at the point (4, 3) is  
 (a)  $x = 3$  (b)  $y = 3$   
 (c)  $x = -3$  (d)  $y = -3$

133.  $x$  and  $y$  are positive number. Let  $g$  and  $a$  be G. M. and AM of these numbers. Also let  $G$  be G. M. of  $x + 1$  and  $y +$

1. If  $G$  and  $g$  are roots of equation

$x^2 - 5x + 6 = 0$ , then

- (a)  $x = 2, y = \frac{3}{4}$  (b)  $x = \frac{3}{4}, y = 12$   
 (c)  $x = \frac{5}{2}, y = \frac{8}{5}$  (d)  $x = y = 2$

134. The co-efficient of  $x^n$  in the expansion of

$$\frac{e^{7x} + e^{3x}}{e^{3x}}$$
 is  
 (a)  $\frac{4n-1 + (-2)^n}{n!}$  (b)  $\frac{4n-1 + 2n}{n!}$

- (c)  $\frac{4n + (-2)^n}{n!}$  (d)  $\frac{4n-1 + (-2)^{n-1}}{n!}$

135. A pair of tangents are drawn from the origin to the circle  $x^2 + y^2 + 20(x + y) + 20 = 0$ , then the equation of the pair of tangent are (a)  $x^2 + y^2 - 5xy = 0$

- (b)  $x^2 + y^2 + 2x + y = 0$   
 (c)  $x^2 + y^2 - xy + 7 = 0$   
 (d)  $2x^2 + 2y^2 + 5xy = 0$

136. If the sum of a certain number of terms of the A.P. 25, 22, 19, ..... is 116. then the last term is  
 (a) 0 (b) 2  
 (c) 4 (d) 6

137. If 1,  $a$  and  $P$  are in A. P. and 1,  $g$  and  $P$  are in G. P.,

then

(a)  $1 + 2a + g^2 = 0$  (b)  $1 + 2a - g^2 = 0$

(c)  $1 - 2a - g^2 = 0$  (d)  $1 - 2a + g^2 = 0$

equal to 138. If  $y = \sin x + e^x$ , then  $dx^2$  is

(a)  $\frac{\sin x e^{-x}}{(\cos x e + x)^2}$  (b)  $\frac{\sin x e^{-x}}{(\cos x e + x)^3}$

$\sin x e + x$

(c)  $\frac{1}{(\cos x e - x)^2}$  (d)  $(-\sin x + e^x)^{-1}$

139. The foci of the hyperbola  $4x^2 - 9y^2 - 1 = 0$  are

(a)  $(\pm \sqrt{13}, 0)$  (b)  $(\pm \sqrt{13}, 0)$

(c)  $(\pm \sqrt{13}, 0)$  (d) None of these

140. From the top of a cliff 50 m high, the angles of depression of the top and bottom of a tower are observed to be  $30^\circ$  and  $45^\circ$ . The height of tower is

(a) 50 m (b)  $50\sqrt{3}$  m  
 (c)  $50(\sqrt{3}-1)$  m (d)  $50(1-\frac{1}{\sqrt{3}})$  m

141. The coefficient of  $x^2$  term in the binomial expansion of  $(x^{1/2} + x^{-1/4})^{10}$  is :

$\frac{1}{10}$

(a)  $\frac{70}{243}$  (b)  $\frac{60}{423}$   
 (c)  $\frac{50}{13}$  (d) none of these

142. The value of  $l$ , for which the circle  $x^2 + y^2 + 2lx + 6y + 1 = 0$  intersects the circle  $x^2 + y^2 + 4x + 2y = 0$  orthogonally, is

(a)  $11/8$  (b)  $-1$   
 (c)  $-5/4$  (d)  $5/2$

143. The value of  $a^2 + b^2 + c^2 + a^2 + b^2 + c^2$  is

(a)  $2(a^2 + b^2 + c^2)$  (b)  $a^2 + b^2 + c^2$   
 (c) 1 (d) None of these

144. If  $f(x) = (a - x^{1/n})^n$ , where  $a > 0$  and  $n \in \mathbb{N}$ , then  $f'(x)$  is equal to :

(a)  $a$  (b)  $x$   
 (c)  $x^n$  (d)  $a^n$

145. Sum of  $n$  terms of the series  $8 + 88 + 888 + \dots$  equals

(a)  $\frac{8}{81} [10^{n+1} - 9n - 10]$

(b)  $\frac{8}{81} [10^n - 9n - 10]$

(c)  $\frac{8}{81} [10^{n+1} - 9n + 10]$

(d) None of these

146. The modulus of the complex number  $z$  such that  $|z + 3 - i| = 1$  and  $\arg(z) = p$  is equal to

(a) 3 (b) 2  
 (c) 9 (d) 4

147. Bag P contains 6 red and 4 blue balls and bag Q contains 5 red and 6 blue balls. A ball is transferred from bag P to bag Q and then a ball

is drawn from bag Q. What is the probability that the ball drawn is blue?

(a)  $\frac{7}{15}$       (b)  $\frac{8}{15}$

(c)  $\frac{4}{19}$       (d)  $\frac{8}{19}$

**148.** The number of 4-digit numbers that can be formed with the digits 1, 2, 3, 4 and 5 in which at least 2 digits are identical, is

(a) 505      (b)  $4^5 - 5!$   
(c) 600      (d) None of these

**149.** Consider the system of linear equations;  $x_1 + 2x_2 + x_3 = 3$   
 $2x_1 + 3x_2 + x_3 = 3$   
 $3x_1 + 5x_2 + 2x_3 = 1$

The system has

- (a) exactly 3 solutions
- (b) a unique solution
- (c) no solution
- (d) infinite solutions

**150.** What is the value of y so that the line through (3, y) and (2, 7) is parallel to the line through (-1, 4) and (0, 6)?

(a) 6      (b) 7  
(c) 5      (d) 9



# SOLUTIONS

## PART - I : PHYSICS

1. (c) The charged sphere is a conductor.

Therefore the field inside is zero and outside it is proportional to  $1/r^2$ .



2. (c) The direction of propagation of electromagnetic wave is perpendicular to

the variation of electric field E as well as to

the magnetic field B.

3. (b) Young's modulus of wire does not vary with dimension of wire. It is a constant quantity.

4. (d) Redraw the given circuit,



$$3R \quad R$$

$$\frac{3R \cdot R}{3R + R} = R^2$$

$$R_{\text{net between AB}} = \frac{3R}{3} + \frac{R}{4R}$$

$$3 \quad 3$$

where,  $R = 16 \text{ W}$

$$R_{\text{net}} = 4 \text{ W}$$

5. (c) According to Kepler's law of planetary motion,  $T^2 \propto R^3$

$$\sqrt{T_2} = \sqrt{\frac{R_2^3}{R_1^3}}$$

$$= 5 \sqrt{\frac{R_2^3}{R_1^3}} = 40 \text{ hours}$$

$$\propto v + v^0 \ddot{o}$$

6. (d)  $f' = f \zeta \frac{v - v_s}{v}$


Here,  $f = 600 \text{ Hz}$ ,  $v_o = 15 \text{ m/s}$

$v_s = 20 \text{ m/s}$ ,  $v = 340 \text{ m/s}$

$$f' = 600 \left( \frac{340 - 15}{340} \right) = 340.340 - 1520 \text{ Hz}$$

$$\sqrt{f'} = 600 \left( \frac{320}{355} \right) \approx 666 \text{ Hz}$$

7. (c) Concave mirror is used as a shaving mirror.



From question :  $v = 15 \text{ cm}$ ,  $u = -10 \text{ cm}$

Radius of curvature,  $R = 2f = ?$


$$\text{Using mirror formula, } \frac{1}{f} + \frac{1}{v} = \frac{1}{R}$$

$$\frac{1}{15} + \frac{1}{(10) - f} = \frac{u}{f}$$

Therefore radius of curvature,  
 $R = 2f = -60 \text{ cm}$

8. (c) From free body diagram,  
 For block to remain stationary,

$$mg \sin a = ma \cos a$$



$$\mu a = g \tan a$$

9. (a) On increasing the temperature, angle of contact decreases.
10. (b) Forward bias opposes the potential barrier and if the applied voltage is more than knee voltage it cancels the potential barrier.

11. (c) In multiplication or division the final result should return as many significant figures as there are in the original number with the least significant figures.

(Rounding off to three significant digits) —

12. (b) Let 'n' be the degree of freedom

$$C_v = \frac{n}{2} R$$

2 also,  $C_p -$

$$C_v = R \quad C_p = C_v + R$$

$$C_p = \frac{n}{2} R + R$$

$$C_p = \frac{5}{2} R$$

so,

$$g = \frac{C_p - C_v}{C_p} = \frac{R}{\frac{5}{2} R} = \frac{2}{5}$$

13. (c) Since range on horizontal plane is

$$R = \frac{u^2 \sin 2\alpha}{g}$$

so it is maximum when,  $\sin 2\alpha = 1$

$$\alpha = 45^\circ$$

14. (c) Mass = 150 gm =  $\frac{150}{1000}$  kg

$$\text{Force} = \text{Mass} \times \text{acceleration}$$

$$= \frac{150}{1000} \times 20 \text{ N} = 3 \text{ N}$$

$$\text{Impulsive force} = F \cdot D = \frac{3}{3} \times 0.103 \text{ N}$$

15. (b) Given,  $d = 10 \text{ m}$

$$q = 30^\circ$$

$$m = 0.5$$

$$F = \sqrt{3} \text{ kN} = \sqrt{3} \times 10^3 \text{ N}$$

$$W = F_s d \cos q$$

Where,  $F_s$

$$= mF$$

$$F_s = 0.5 \times \sqrt{3} \text{ kN}$$

$$F_s = 0.866 \text{ kN}$$

$$F_s = 866 \text{ N}$$

$$\text{So, } W = 866 \times 10 \times \cos 30^\circ$$

$$W = \frac{866 \times 10 \times \sqrt{3}}{2}$$

$$W = 7499.56 \text{ J}$$

$$W \approx 7.5 \text{ kJ}$$

$$\vec{r} = (2\hat{i} + 3\hat{j} + 4\hat{k}) (3\hat{i} + 4\hat{j} + 5\hat{k})$$

16. (a)  $W = F \cdot s$

$$= 2 \times 3 + 3 \times 4 + 4 \times 5 = 38 \text{ J}$$

$$P = \frac{W}{t} = \frac{38}{4} = 9.5 \text{ W}$$

17. (d)  $v_e = \sqrt{\frac{2GM}{R}}$

$$\text{and, } v_g = \sqrt{\frac{2GM}{R}} = \sqrt{\frac{2GM}{R}} = \frac{v_e}{\sqrt{2}}$$

$$\frac{v_g}{v_e} = \frac{1}{\sqrt{2}}$$

18. (b)  $\frac{dA}{dt} = \frac{L}{2m} = \text{Constant}$

19. (a) According to Bernoulli's theorem

$$P_1 + \frac{1}{2} \rho v_1^2 = P_2 + \frac{1}{2} \rho v_2^2 \dots (i)$$

According to the condition,

$$P_1 - P_2 = \frac{1}{2} \rho (v_2^2 - v_1^2) = 5$$

From equation of continuity,

$$A_1 v_1 = A_2 v_2 \text{ so, } A_1^2 v_1^2 = A_2^2 v_2^2$$

$$5 \rho v_2 = 5 v_1$$

From equation (i)

$$P_1 - P_2 = \frac{1}{2} \rho (v_2^2 - v_1^2)$$

2 -

$$f' = \frac{1}{2p} \sqrt{\frac{4k_1 + 4k_2}{m}}$$

$$f' = \frac{1}{2p} \sqrt{\frac{4(k_1 + k_2)}{m}} = 2 \times \frac{1}{2p} \sqrt{\frac{k_1 + k_2}{m}} = 2f$$


$$\text{or } 3 \times 10^5 = 12 \sqrt{1000 \cdot 25 (v_1^2 - v_1^2)}$$

$$p \cdot 600 = 24 \sqrt{v_1^2 - v_1^2} = 25$$

$$\sqrt{v_1^2 - v_1^2} = 5 \text{ m/s}$$

20. (c) Work done = Area under curve ACBDA

21. (a) B



From figure,  $V_B \sin \theta = V_w$

$$\sin \theta = \frac{V_w}{V_B} = \frac{1}{2} \Rightarrow \theta = 30^\circ \quad [\because V_B = 2V_w]$$

Time taken to cross the river.

$$t = \frac{D}{V \cos \theta} = \frac{D}{V_B \cos 30^\circ} = \frac{2D}{V_B \cos 30^\circ}$$

22. (a) The two springs are in parallel. Effective spring constant,  $k = k_1 + k_2$

Now, frequency of oscillation is given by

$$f =$$

or,  $f = \dots(i)$

When both  $k_1$  and  $k_2$  are made four times their original values, the new frequency is given by

$$A = A_0 e^{-bt/2m}$$

Case 1 :-

$$A^0$$

$$\text{When } t = 2 \text{ s, } A = \frac{A_0}{3}$$

23. (b) Drift velocity,  $v_d = \frac{I}{n e A} = \frac{J}{n e} = \frac{E}{V} = \frac{1}{3} V$

$$v_d = \frac{I}{n e A} = \frac{J}{n e} = \frac{E}{V} = \frac{1}{3} V$$

24. (b) Amplitude of a damped oscillator

$$A_3 = A_0 e^{-2b/2m} = e^{-b/m} \dots (i)$$

Case 2 :-

$$A^0$$

$$\text{When } t = 6 \text{ s, } A = \frac{n}{3}$$

$$A_6 = A_0 e^{-6b/2m} = (e^{-b/m})^3 \dots (ii)$$

From (i) and (ii)

$$\ln \frac{n}{3} = -3 \frac{b}{m} \Rightarrow \frac{b}{m} = \frac{1}{3} \ln \frac{n}{3}$$

25. (d) Angular speed of electron in the nth orbit of Bohr H-atom is inversely proportional to  $n^3$

$$\omega \propto \frac{1}{n^3}$$

26. (a) C = equivalent capacitance

$$\frac{1}{C} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C = 1 \text{ mF}$$

$$C_1 = 2 \text{ mF}, C_2 = 3 \text{ mF}, C_3 = 6 \text{ mF}$$

Charge on each capacitor in series circuit will be same.

$q = CV = (1 \times 10^{-6}) \times 10 = 10 \text{ mC}$   
 Charge across  $3 \mu\text{F}$  capacitor will be  $10 \mu\text{C}$ .  
 E

27. (a)  $E_{12} = \frac{1}{4\pi\epsilon_0} \left[ \frac{q}{r^2} - \frac{q'}{(2x)^2} \right]$   
 $\frac{1}{4\pi\epsilon_0} \left[ \frac{(500)(600)}{44^2} - \frac{(300)(300)}{44^2} \right] = \dots$

28. (c) v

29. (a) For open pipe,  $n_1 = \frac{v}{2\ell}$ , where  $n_1$  is the fundamental frequency of open pipe. length of open pipe is,

$\ell = \frac{2v}{n} = \frac{2 \times 330}{300} = 1120$

$\propto v$

1st overtone of open pipe,  $n_2 = 2n_1 = 2 \times 300 = 600$

$\frac{2\ell}{\lambda} = 2$  Ist overtone of closed pipe,  $\propto v$

$n_3 = 3n_1 = 3 \times 300 = 900$   $\frac{4\ell'}{\lambda} = 3$  where,  $\ell'$  = length of closed pipe As freq. of 1st overtone of open pipe = freq.

of 1st overtone of closed pipe

$\frac{v}{\lambda} = \frac{v}{\lambda} \Rightarrow \frac{3\ell}{3} = \frac{4\ell'}{4} \Rightarrow \ell' = \frac{3}{4}\ell = \frac{3}{4} \times 1120 = 840 \text{ cm}$

$\frac{2\ell}{\lambda} = 2 \Rightarrow \ell = \frac{\lambda}{2} = \frac{300}{2} = 150 \text{ cm}$

$D_1(2)D_1 = 4b$

30. (d)  $b = \frac{d}{2}$  and  $b\phi = (d/2)$

d

Thus the fringe width becomes four times.

31. (a) As we know,  $t$  is change in angular momentum.

$\frac{20}{3}$

so,  $t = \frac{20}{3}$  SI units

32. (a) Let  $q$  and  $q'$  be the charges on spheres of radii  $x$  and  $2x$  respectively.

Given,  $q + q' = Q$  ... (i) Surface charge densities are

$s = \frac{q}{4\pi x^2}$  and  $s' = \frac{q'}{4\pi (2x)^2}$

Given,  $s = s'$

$\frac{q}{4\pi x^2} = \frac{q'}{4\pi (2x)^2}$


$q' = 4q$

From eq. (i),  $q' = Q - q$  or,  $4q = Q - q$

or,  $Q = 5q$  ... (ii)

$q' = Q - q = Q - \frac{Q}{5} = \frac{4Q}{5}$

33. (d)  $L = 2 \text{ H}, E = 5 \text{ volts}, R = 1 \text{ W}$



$\frac{E}{Z}$

$\sqrt{R^2 + (\omega L)^2}$

5

$\sqrt{R^2 + (\omega L)^2}$

5

$\frac{5}{\sqrt{1 + (200\pi)^2}} = \frac{5}{200\pi}$

I =

Z =

I =

Energy in inductor =  $\frac{1}{2} LI^2$

$= \frac{1}{2} \times 2 \times \left(\frac{5}{200\pi}\right)^2$

$= 6.33 \times 10^{-5} \text{ joules}$

34. (b)  $F = Bi\ell = 2 \times 1.2 \times 0.5 = 1.2 \text{ N}$

35. (b) Distance between two cars leaving from the station A is,

$d = \frac{1}{6} \times 60 = 10 \text{ km}$

Man meets the first car after time,

$$t_1 = \frac{60}{60 + 60} = \frac{1}{2} \text{ h}$$

He will meet the next car after time,

$$t_2 = \frac{60}{10} + \frac{60}{60} = 12 \text{ h}$$

In the remaining half an hour, the number

$$\text{of cars he will meet again is, } n = \frac{1/2}{12} = 6$$

\ Total number of cars would be meet on route will be 7.

36. (b) From  $v = r\omega$ , linear velocities ( $v$ ) for particles at different distances ( $r$ ) from the axis of rotation are different.

37. (a) For an SHM, the acceleration  $a = -\omega^2 x$

$$\text{where, } \omega \text{ is a constant} = \frac{2\pi}{T}$$

$$a = -\left(\frac{2\pi}{T}\right)^2 \cdot x$$

The period of oscillation  $T$  is a constant.

$$\frac{a}{T^2} \text{ is a constant. } x$$


38. (a) As the inward magnetic field increases, its flux also increases into the page and so induced current in bigger loop will be anticlockwise. i.e., from D to C in bigger loop and then from B to A in smaller loop.

39. (a) Since A moves twice the distance moved by B.

If acceleration of B is 'a', then acceleration of A is '2a'.

$T' - (T + T) = 0$  (since pulley is massless)

$$\text{p } T' = 2T \quad \dots(i)$$



For 5 kg block

$$5g - T' = 5a$$

for 2 kg block

$$\text{p } 5g - 2T = 5a$$

$$T = 2 \times (2a) = 4a$$

From equations (ii) and (iii),

$$5g - (2 \times 4a) = 5a$$

$$5g - 8a = 5a$$

$$5g = 13a$$

$$a = \frac{5g}{13}$$

$$13 a_A = 2a \quad 10 \text{ ---}$$

$$13g ; a_B = a \quad \text{---} 513g$$

40. (a) Isotones means equal number of neutrons  
i.e.,  $(A-Z) = 74 - 34 = 71 - 31 = 40$

43. (b)

41. (a)  
42. (c)

As per Arrhenius equation ( $k = Ae^{-E/RT}$ ), the rate constant increases exponentially with temperature.

Mass of O<sub>2</sub> absorbed per gram of adsorbent

$$= \frac{3.6}{1.2} = 3$$

$$1.2$$

No. of moles of O<sub>2</sub> absorbed per gram of

$$\text{adsorbent} = \frac{3}{32}$$

Volume of O<sub>2</sub> absorbed per gram of adsorbent


$$PV = nRT$$

$$nRT$$

$$V = \frac{nRT}{P}$$


$$32 \times \frac{3}{32} = 2.1$$

$$= 3 \times 0.0821 \times 273$$




(Impure)


Volatile compound





Thermal decomposition pure



I<sub>2</sub> gives violet colouration in CHCl<sub>3</sub>.



Clathrate formation involves dipole-induced dipole interaction. Eu (63) = [Xe]




47. (a) As positive charge on the central metal


can donate electron density into the p\* orbitals of CO ligand (donation of electron density into p\* orbitals of CO result in weakening of C – O bond). Hence, the C –


O bond would be strongest in [Mn(CO)<sub>6</sub>]<sup>+</sup>.

atom increases, the less readily the metal




Four monochloro derivatives are chiral.







51. (b) In compound (b), the lone pair of nitrogen is not involved in resonance therefore it is the strongest base.



52. (c) Sucrose, being a non-reducing sugar, does not reduce Benedict's solution. Remember that fructose has an α-hydroxy ketonic group, which is also reducing group (different from ordinary ketonic group)



53. (c) Zn<sup>2+</sup> present in alternate tetrahedral void






$$\frac{1}{2} = 0.5$$

S<sup>2-</sup> present in ccp = 4


(Major)  
Z  
Now, during electrophoresis, glycine moves towards anode.

Zn<sub>4</sub>S<sub>4</sub> = ZnS i.e., AB type compound.

54. (d) Glycine in alkaline solution exists as anion and migrates to anode.



Due to internal proton transfer of H<sup>+</sup> from the -COOH group to the -NH<sub>2</sub>, the amino acid exists as an ion with both a negative charge and a positive charge, called a Zwitter ion



$$\text{Ka} = \frac{[\text{H}^+][\text{X}^-]}{[\text{HX}]}$$

Adding an alkali to glycine



$$10^{-5} = \frac{[\text{H}^+] \cdot 0.05}{\text{_____}}$$






$$\frac{0.20}{0.20005 - 5} + ] \cdot 10 = [H$$

$$4 \times 10^{-5} M = [H^+]$$

57. (c) Pt | H<sub>2</sub> (640 mm) | HCl | H<sub>2</sub> (510 mm) | Pt



$$E^\circ_{\text{cell}} = 0$$





$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.0591}{n} \log \frac{p_2}{p_1}$$

$$= 0 - 0.0591 \log \frac{510}{640}$$

$$= 2.91 \times 10^{-3} V$$



58. (a)



XeF<sub>4</sub> has zero net dipole moment




59. (a) Boron has the highest ionisation enthalpy amongst the following. Ionisation enthalpy decreases down the group and increases across the period.

60. (a) Element with atomic number 7 has the smallest size and highest ionization enthalpy Nitrogen – Atomic Number 7



N has a stable half-filled electronic configuration therefore it is difficult to remove electron and hence it has a high ionization enthalpy.

61. (a) Except dacron all are additive polymers. Terephthalic acid condenses with ethylene glycol to give dacron.



62. (a) Phenelzine is an antidepressant, while others are antacids.


63. (b) 1 molal solution means 1 mole of solute dissolved in 1000 g solvent.

$$\backslash \quad n_{\text{solute}} = 1 \quad w_{\text{solvent}} = 1000 \text{ g}$$

$$\backslash \quad n_{\text{solvent}} = \frac{1000}{18} = 55.56 \quad x_{\text{solute}} =$$

$$\frac{1}{1 + 55.56} = 0.0177$$

64. (a)  $\text{ClCH}_2\text{CH}_3$



trans-2-Chloro-3-iodo-2-pentene

65. (a) Stability of carbocation  $\mu$  no. of  $\alpha$ -H

present on carbocation.

66. (a) Only Lindlar's catalyst converts alkyne to

alkene (*cis* addition) and alkenes with

Baeyer's reagent give *cis* glycols.

67. (b) Lower oxidation state become more stable

on moving down the group

$\text{Al} < \text{Ga} < \text{In} < \text{Tl}$

68. (a)  $\text{Be}(\text{OH})_2$  is amphoteric while  $\text{Ca}(\text{OH})_2$ ,


$\text{Sr}(\text{OH})_2$  and  $\text{Ba}(\text{OH})_2$  are all basic.

69. (a)  $\text{H}_2\text{O}_2 + 2\text{KI} \rightarrow 2\text{I}_2 + 2\text{H}_2\text{O}$ , O.S. of I (-1) changes

to  $\text{I}_2$  (Zero) There is increase in oxidation

number, hence oxidation.

70. (d) The balanced equation is



71. (a)  $[\text{A}] = 2$ ,  $[\text{B}] = 4$  and  $[\text{C}] = 6$

Multiply the three equations,

$$2 \times 4 \times 6 = [\text{A}] = 48$$

72. (d)  $\text{DG} = \text{DH} - \text{TDS} = +ve$

for spontaneous change,  $\text{DH} < 0$ ,  $\text{DS} > 0$  for non-spontaneous change,  $\text{DH} > 0$ ,  $\text{DS} < 0$

73. (b)  $r_1^2 \sqrt{\frac{d_2}{d_1}} = \sqrt{\frac{16}{1}} \implies 4:1$

74. (a)  $2\text{PCl}_5 \rightleftharpoons \text{PCl}_4^+ + \text{PCl}_6^-$

75. (a) Isoelectronic species have same no. of electrons.

Ions	$\text{O}^{2-}$	$\text{F}^-$	$\text{Na}^+$
$\text{Mg}^{2+}$			
$8 + 2$	$9 + 1$	$11 - 1$	$12 - 2$
No. of $e^- = 10$	$10$	$10$	$10$

$\text{O}^{2-}, \text{F}^-, \text{Na}^+, \text{Mg}^{2+}$  are isoelectronic


76. (c) This is Avogadro's hypothesis. According to this, equal volume of all gases contain equal no. of molecules under similar condition of temperature and pressure. 77. (a) Eq. of A = Eq. of B

$$mE_A = mE_B; E_A = mm' E_B$$


A B

78. (c) For  $4p$  electron  $n = 4$ ,  $l = 1$ ,  $m = -1, 0, +1$  and  $s = +\frac{1}{2}$  or  $-\frac{1}{2}$

79. (c)  $l = 2$  represent  $d$  orbital for which



80. (a)  $\text{B}_2$  is paramagnetic due to the presence of unpaired electron in  $p_x$  or  $p_y$  orbital.



M.O diagram for  $\text{B}_2$  molecule


**PART - III (A): ENGLISH  
PROFICIENCY**

81. (a) The word **Garrulous** (Adjective) means : talkative; talking a lot.
82. (b) The word **Tinsel** (Noun/Adjective) means :  
strips of shiny material like metal used as decorations.
83. (a) The word **Labyrinth** (Noun) means : a place that has many confusing paths or passage. The correct synonym will be 'meandering' which means, 'to have a lot of curves on a path'.
84. (b) Knack means a clever way of doing something.
85. (d) Pernicious means highly injurious or destructive.
86. (d) Opulence means wealthy.
87. (a) In good friendships, we receive as much as we give.
88. (b) Empathy means the ability to show and understand the feelings of others.
89. (c) A strong friendship helps us gain acceptance and tolerance.
90. (d) The very first line of the passage states that friendships and relationships grow when they are nurtured just like nurturing a plant.
91. (a) **Dependent on** = needing somebody / something in order to survive or be successful; affected or decided by something.
92. (b) **Take your leave** = to say good bye.
93. (b) Here, indefinite article i.e., 'about a plane crash' should be used. No particular incident is evident here.
94. (b) 'With a View to' should be followed by gerund i.e., surveying.
95. (a) Here, time period is given. Hence, Past Perfect Continuous i.e., 'It had been lying' ....should be used.

**PART - III (B) : LOGICAL  
REASONING**

96. (b) 97. (c) 98. (d)

99. (c) Medicine is given to patient. Similarly, Education is given to student.




101. (a) Except 5, all numbers are perfect square numbers.

102. (a) As,  
TIRED = 20 + 9 + 18 + 5 + 4 = 56  
BRAIN = 2 + 18 + 1 + 9 + 14 = 44  
Similarly,  
LAZY = 12 + 1 + 26 + 25 = 64.

103. (b)  $8 \times 8 \times 88 = 5632$   $7 \times 7 \times 77 = 3773$   
Similarly,  $6 \times 6 \times ? = 3132$

3132

$$\sqrt{?} = \frac{3132}{66} = \boxed{87}$$



105. (d) Mohan is son of Ram Lal and uncle of Ram and Rekha. Mithun is uncle of Sharat who is son of Rekha. Rekha is niece of Mohan. Therefore, Mithun is brother of Rekha's husband.

**PART - IV : MATHEMATICS**

106. (c) Given line is  $x \cos a + y \sin a = P$  ....(1)

Any tangent to the ellipse is  $x \cos \theta + y \sin \theta = p$

$$P = p \quad \dots(2) \quad a \quad b$$

Comparing (1) and (2)

$$\cos \theta = \cos a \quad \sin \theta = \sin a$$

$$a \cos \alpha \quad b \sin \alpha \quad P$$

$$\frac{P}{b \sin \alpha} \cos \theta = \frac{a \cos \alpha}{P} \text{ and } \sin \theta = \frac{P}{a \cos \alpha}$$

Eliminate  $q$ ,  $\cos^2 q + \sin^2 q$

$$= \frac{a \cos^2 \alpha}{P^2} + \frac{b \sin^2 \alpha}{P^2}$$

$$\text{or } a^2 \cos^2 \alpha + b^2 \sin^2 \alpha = P^2$$

107. (c) As  $a_1, a_2, a_3, \dots, a_n$  are in A.P. we get,  $a_2 - a_1 = a_3 - a_2 = \dots = a_n - a_{n-1} = d$  (say)

$$\text{Now, } \frac{1}{\sqrt{a_1} + \sqrt{a_2}} = \frac{\sqrt{a_1} - \sqrt{a_2}}{a_1 - a_2} = \frac{\sqrt{a_1} - \sqrt{a_2}}{-d}$$

Similarly,

$$\frac{1}{\sqrt{a_2} + \sqrt{a_3}} = \frac{\sqrt{a_2} - \sqrt{a_3}}{-d}, \dots, \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} = \frac{\sqrt{a_{n-1}} - \sqrt{a_n}}{-d}$$

$$\therefore \frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} = \frac{\sqrt{a_1} - \sqrt{a_2} + \sqrt{a_2} - \sqrt{a_3} + \dots + \sqrt{a_{n-1}} - \sqrt{a_n}}{-d}$$

$$= \frac{\sqrt{a_1} - \sqrt{a_n}}{-d} = -\frac{1}{d} \left[ \frac{a_1 - a_n}{\sqrt{a_1} + \sqrt{a_n}} \right]$$

$$= -\frac{1}{d} \left[ \frac{a_1 - \{a_1 + (n-1)d\}}{\sqrt{a_1} + \sqrt{a_n}} \right]$$

[formula for  $n^{\text{th}}$  term]

$$= -\frac{1}{d} \frac{-(n-1)d}{\sqrt{a_1} + \sqrt{a_n}} = \frac{n-1}{\sqrt{a_1} + \sqrt{a_n}}$$

108. (b) Given differential equation is :  $x \cos x \frac{dy}{dx} + y(x \sin x + \cos x) = 1$  Dividing both the sides by  $x \cos x$ ,

$$P \frac{dy}{dx} + y \frac{\sin x + \cos x}{\cos x} = \frac{1}{x \cos x}$$

$$P \frac{dy}{dx} + y \tan x + \frac{y}{x \cos x} = \frac{1}{x \cos x}$$

$$P \frac{dy}{dx} + y \sec x \tan x + \frac{y}{x} = \frac{1}{x \cos x}$$

$$\frac{dy}{dx} + y \sec x \tan x + \frac{y}{x} = \frac{1}{x \cos x}$$

which is of the form  $\frac{dy}{dx} + Py = Q$

$$\text{Here, } P = \frac{\tan x}{x} \text{ and } Q = \frac{1}{x \cos x} = \frac{\sec x}{x}$$

$$\text{Integrating factor} = e^{\int P dx} = e^{\int \frac{\tan x}{x} dx} = e^{\log \sec x} = \sec x$$

$$= e^{(\log \sec x + \log x)} = e^{\log(\sec x \cdot x)} = x \sec x$$

109. (c) Equation of the first line  $L_1$  is

$$x - 1 = y - 2 = z - 3 \text{ and that of the second line } 2x - 3 = y - 4 = z - 5$$

$L_2$  is  $x - 5 = y - 4 = z - 0$  Clearly, these lines are not parallel (the ratios of D.R. are not equal). Any point P on the first line is  $(1 + 2l, 2 + 3l, 3 + 4l)$  and any point Q on the second line is  $(4 + 5m, 1 + 2m, m)$ . If these two points P and Q are identical then.

$$1 + 2l = 4 + 5m \dots (1)$$

$$2 + 3l = 1 + 2m \dots (2)$$

$$3 + 4l = m \dots (3)$$

From (2) and (3), we get  $l = m = -1$ , which also satisfies (1). Thus the two lines  $L_1$  and  $L_2$  intersect and the coordinates of the point of intersection are  $(-1, -1, -1)$ .

110. (c) We have  $y - x \frac{dy}{dx} = a y^2 + \frac{dy}{dx}$

$$P y dx - x dy = a y^2 dx + dy$$

$$\int y(1-ay)dx = \int (x+a)dy$$

dy

$$\int x+a - y(1-ay) = 0$$

Integrating, we get


$$\log(x+a) - \log y + \log(1-ay) = \log C$$

$$\log \frac{(x+a)(1-ay)}{y} = \log C \text{ i.e. } (x+a)(1-ay) = Cy$$

Since the curve passes through (0, 1) and (1, 0)

$$\log \frac{(0+a)(1-1)}{1} = \log C \text{ i.e. } C = -4a^2$$

$$\text{So, } (x+a)(1-ay) = -4a^2y$$



111. (c)

Equation of given circle is  $x^2 + y^2 = 4$

Its centre,  $O = (0, 0)$  and radius,  $r = 2$

Draw  $OM \perp AB$

Clearly M is the mid-point of AB which subtends a right angle at O.

In  $\triangle OAB$ ,  $OA = OB$  radius

$$\angle OAB = \angle OBA = A$$

4 and in

$$\triangle OMA, \sin A = \frac{OM}{OA}$$

$$\frac{OM}{2}$$

$$\sin A = \frac{OM}{2}$$

$$\frac{OM}{2} = \sin A$$

$$OM = 2 \sin A \quad \dots(1)$$

$$\text{Let } M = (x, y) \text{ then } OM = \sqrt{x^2 + y^2} \quad \dots(2)$$

From (1) and (2),  $x^2 + y^2 = 2 \sin^2 A$  This is the required equation of locus.

112. (c)  $\int \frac{1}{1 + \sin A - \cos A} dx$

$$\begin{aligned} \int \frac{1}{1 + \sin A - \cos A} dx &= \int \frac{1 + \cos A}{1 + \sin A - \cos A} dx \\ &= \int \frac{2 \cos^2 \frac{A}{2}}{2 \sin^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2}} dx \\ &= \int \frac{2 \cos^2 \frac{A}{2}}{2 \cos \frac{A}{2} (\sin \frac{A}{2} + \cos \frac{A}{2})} dx \\ &= \int \frac{\cos \frac{A}{2}}{\sin \frac{A}{2} + \cos \frac{A}{2}} dx \end{aligned}$$

113. (c)

$$\frac{1}{1 + \sin A + \cos A}$$

$$\begin{aligned} &\frac{2 \sin^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2}}{2 \cos^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2}} \\ &= \frac{2 \cos^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2}}{2 \cos^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2}} \end{aligned}$$

$$= \frac{2 \cos^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2}}{2 \cos^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2}} = \tan A$$

$$2 \cos^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2} = 2 \cos \frac{A}{2} (\cos \frac{A}{2} + \sin \frac{A}{2})$$

114. (c) Given  $\sqrt{1+x} + y\sqrt{1+y} = 0$

$$\sqrt{1+x} = -y\sqrt{1+y}$$

Squaring both sides, we get  $x^2(1+y)$

$$= y^2(1+x)$$

$$x^2 - y^2 + x^2y - xy^2 = 0 \Rightarrow (x-y)(x+y+xy) = 0$$

$$p y = x \text{ or } y(1+x) = -x \quad p y = x \text{ or } y = \frac{-x}{1+x}$$

1 + x

$$p dy = \frac{-x(1+x) + x \cdot 1}{(1+x)^2} = \frac{-x}{(1+x)^2}$$

115. (b) Given :  $f(x) = 3x^4 + 4x^3 - 12x^2 + 12$   
Differentiating with respect to x, we get  $f'(x) = 12x^3 + 12x^2 - 24x$

For f(x) to be increasing  $f'(x) > 0$


$$0 < 12x^3 + 12x^2 - 24x > 0$$

$$p 12x(x^2 + x - 2) > 0$$

$$p 12x(x-1)(x+2) > 0$$

$$p x(x-1)(x+2) > 0$$

$$p -2 < x < 0 \quad \text{or} \quad x > 1$$



It means  $x \in (-2, 0) \cup (1, \infty)$ .

Hence f(x) is increasing in  $(-2, 0)$  and  $(1, \infty)$

116. (c) Consider  $x^2 + y^2 = 1$ ,  $x^2 + y^2 = 4$

$x, y \geq 0$  convert them into equation and solve them and draw the graph of these equations we get  $y = 1$  and  $x = 3/2$

117. (b) Here,  $a = 2\hat{i} + 5\hat{j} - 3\hat{k}$ ,  $N = 6\hat{i} - 3\hat{j} + 2\hat{k}$  and  $d = 4$ .

Therefore, the distance of the point  $(2, 5, -3)$  from the given plane is

$$\frac{|(2\hat{i} + 5\hat{j} - 3\hat{k}) \cdot (6\hat{i} - 3\hat{j} + 2\hat{k}) - 4|}{\sqrt{6^2 + 3^2 + 2^2}} = \frac{|12 - 15 - 6 - 4|}{7} = \frac{13}{7}$$

distance =  $\frac{13}{7}$

118. (a)  $I = \int_0^{\pi/2} \log(\tan x) dx = \int_0^{\pi/2} \log \tan x dx - \int_{\pi/2}^0 \log \tan x dx$

$$p 2 = \int_0^{\pi/2} \log \cot x dx$$

$$x dx \left( \int_0^{\pi/2} \log \cot x dx \right)$$

$$\int_0^{\pi/2} 2I = \int_0^{\pi/2} \log \tan x dx + \int_0^{\pi/2} \log \cot x dx$$

$$0 \int_0^{\pi/2} \log \tan x dx + \int_0^{\pi/2} \log \cot x dx$$

$$= \int_0^{\pi/2} [\log \tan x + \log \cot x] dx$$

$$= \int_0^{\pi/2} \log(\tan x \cdot \cot x) dx = \int_0^{\pi/2} \log 1 dx = 0$$

119. (a)

120. (c) Since,  $(e^x + 1) y dy = (y + 1)e^x dx$

$$p \frac{dy}{y} = \frac{dx}{1+y} + \frac{dx}{(1+y)e^x}$$

$$dx \approx y \approx e^x + 1$$

$$p dy = \frac{dx}{1+y} + \frac{dx}{(1+y)e^x}$$

$$\approx y \approx e^x + 1$$

$$p \int \frac{1}{1+y} dy = \int \frac{1}{1+y} dy$$

$$= \int \frac{1}{1+y} dy$$

After integrating on both sides, we have

$$\int \frac{1}{1+ye^{xy}} dy = \int \frac{1}{1+e^{xy}} dx$$

$$\int \frac{1}{1+ye^{xy}} dy = \int \frac{1}{1+e^{xy}} dx$$

$\int y - \log |(1+y)| = \log |(1+e^x)| + \log k$   
 Hence  $y = \log [k(1+y)(1+e^x)]$

121. (c) Equation of parabola is  $y^2 = 4ax$

$$\frac{dy}{dx} = \frac{2a}{y} \quad [\text{slope of tangent}] dx$$

$\frac{dy}{dx} = \frac{2a}{y}$  (On differentiating w.r.t 'x')

So, slope of normal =  $-\frac{y}{2a}$  (at  $(2at^2, 2at)$ )

$$= -\frac{2at}{2a} = -t$$

$$\frac{p}{2}$$

122. (a) Let  $I = \int_0^{\frac{\pi}{2}} x \sin^2 x \cos^2 x dx \quad \dots(i)$

From the definite integral property

$$\int_0^a f(x) dx = \int_0^a f(a-x) dx$$

we have

$$\frac{p}{2}$$

$$I = \int_0^{\frac{\pi}{2}} \frac{p}{2} - x \sin^2 x \cos^2 x dx$$

... (ii)  $\int_0^{\frac{\pi}{2}} \frac{p}{2} - x \sin^2 x \cos^2 x dx$

$$\cos^2 x = \frac{1 + \cos 2x}{2}$$

$$\sin^2 x = \frac{1 - \cos 2x}{2}$$

By adding (i) and (ii)

$$\frac{p}{2}$$

$$\int_0^{\frac{\pi}{2}} \frac{p}{2} \cos^2 x dx = 2I$$

$$\text{or } 2I = \int_0^{\frac{\pi}{2}} 2 \sin 2x dx$$

$$[\because \sin 2x = 2 \sin x \cos x]$$

$$= \int_0^{\frac{\pi}{2}} 2(1 - \cos 4x) dx \quad (\because \cos 2q = -12 \sin 2q)$$

$$\frac{p}{2} - \frac{\sin 4x}{4}$$

$$\int_0^{\frac{\pi}{2}} \frac{p}{2} - \frac{\sin 4x}{4}$$

$$\int_0^{\frac{\pi}{2}} \frac{p}{2} - \frac{\sin 4x}{4} = \frac{p}{2} \left[ \frac{x}{1} - \frac{-\cos 4x}{4} \right]_0^{\frac{\pi}{2}}$$

$$\left| \begin{array}{cc|c} 1 & & \\ 3i & -1 & \\ & i & \end{array} \right| \begin{array}{c} 6i - 3i \\ \mathbf{123. (d)} \end{array}$$

$$20 \quad 3$$

$$= 6i [3i^2 + 3] + 3i [4i + 20] + 1 [12 - 60i]$$

$$= 6i [-3 + 3] + 12i^2 + 60i + 12 - 60i$$

$$= -12 + 12 = 0 = x + iy$$

$$\setminus x = 0$$

3

129. (b) From the given problem:  $P(A \dot{\cup} B) =$

$\rightarrow$

4

124. (a)  $\lim_{x \rightarrow 0} \frac{\log_e(1+x) + x^2 - x}{x^2}$

$$= \lim_{x \rightarrow 0} \frac{\log_e(1+x) + x^2 - x}{x^2}$$

$$= \lim_{x \rightarrow 0} \frac{x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \dots + x^2 - x}{x^2}$$

125. (a)  $2 \cos^2 x + 3 \sin x - 3 = 0$   
 $2 - 2 \sin^2 x + 3 \sin x - 3 = 0 \Rightarrow$   
 $\sin x - 1)(\sin x - 1) = 0$

$$\Rightarrow \sin x = \frac{1}{2} \text{ or } \sin x = 1$$

$$\Rightarrow x = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{2\pi}{3}, \text{ i.e. } 30^\circ, 150^\circ, 90^\circ.$$

126. (b)

127. (d)  $y = \tan^{-1} \frac{\sqrt{x}}{1+x} - \tan^{-1} \frac{\sqrt{x}}{1+x^2}$

$$= \tan^{-1} \left( \frac{x}{1+x} \right) - \tan^{-1} \left( \frac{x}{1+x^2} \right)$$

On differentiating w.r.t. x, we get

$$y' = \frac{1}{1+x} \cdot \frac{1}{1+x} - \frac{1}{1+x^2} \cdot \frac{1}{1+x^2}$$

$$\Rightarrow y'(1) = \frac{1}{2} \cdot \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$$

128. (b) The diagonal = R Thus the area of rectangle

$$= \frac{1}{2} R^2 = \frac{R^2}{2}$$

$$P(A \cap B) = \frac{1}{4}$$

$$\Rightarrow \frac{1}{4} = 2 - P(A) - P(B) \Rightarrow P(A) = \frac{3}{4}$$

$$\frac{1}{2} P(A)$$

3

3

$$P(A \dot{\cup} B) = P(A) + P(B) - P(A \cap B)$$

$$\Rightarrow P(B) = P(A \dot{\cup} B) + P(A \cap B) - P(A)$$

$$= \frac{3}{4} + \frac{1}{4} - \frac{3}{4} = \frac{1}{4}$$

$$\frac{e^{1/x} - e^{-1/x}}{e^{1/x} + e^{-1/x}}$$

130. (b)  $\lim_{x \rightarrow 0} \frac{e^{1/x} + e^{-1/x}}{e^{1/x} - e^{-1/x}}$

$$\lim_{x \rightarrow 0} \frac{e^{1/x} + e^{-1/x}}{e^{1/x} - e^{-1/x}}$$

$$= \lim_{x \rightarrow 0} \frac{1 + e^{-2/x}}{1 - e^{-2/x}} = +\infty$$

$$\lim_{x \rightarrow 0} \frac{e^{1/x} - e^{-1/x}}{e^{1/x} + e^{-1/x}} = \lim_{x \rightarrow 0} \frac{e^{1/x}(e^{2/x} - 1)}{e^{1/x}(e^{2/x} + 1)}$$

$$= \lim_{x \rightarrow 0} \frac{e^{1/x} \cdot 2e^{2/x}}{e^{1/x} \cdot 2e^{2/x}} = 1$$

$$= \lim_{x \rightarrow 0} \frac{2e^{3/x}}{2e^{3/x}} = 1$$

Limit doesn't exist, so f(x) is not continuous at 0.

131. (b) Put  $x = \cos 2q$

$$\int \cos \{2 \tan^{-1} \sqrt{2 \sin 2q} - q\} dq$$



$$= -\frac{1}{4} \int \frac{\sin 4q}{\cos 4q + c} dq$$

$$= -\frac{1}{4} \int \frac{1 - \tan^2 2q}{1 + \tan^2 2q} \cdot 2 \tan 2q \cdot 2 dq$$

$$= -\frac{1}{2} \int \frac{1 - t^2}{1 + t^2} \cdot 2t dt$$

$$= -\frac{1}{2} \int \left( \frac{1 - t^2}{1 + t^2} \right) \cdot 2t dt$$

132. (b)  $T = S_1 \Rightarrow x(4) + y(3) - 4(x+4) = 16 + 9 - 32 \Rightarrow 3y - 9 = 0 \Rightarrow y = 3$

133. (d) The roots of equation are 2 and 3

$$\sqrt{g} = \sqrt{xy} = 2 \Rightarrow xy = 4$$

$$G = \sqrt{(x+1)(y+1)} = 3 \Rightarrow (x+1)(y+1) = 9$$

$$\Rightarrow x = y = 2$$

134. (c)  $\frac{e^{7x} e^{3+x}}{e^{-2x}} = e^{4x+10}$

$$= e^{4n} \cdot \frac{(4x)^2}{n!} + \dots + e^{1+4x} + \dots + e^{1-(2x)} + \dots + e^{2!} + \dots + e^{2!} + \dots$$

$$\backslash \text{coeff. of } x^n = \frac{4^n}{n!} + \frac{(-2)^n}{n!}$$

135. (d) Equation of pair of tangents is given by  $SS_1 = T^2$ ,

$$\text{or } S = x^2 + y^2 + 20(x+y) + 20, S_1 = 20, T = 10(x+y) + 20 = 0$$

$$\backslash SS_1 = T^2$$

$$\Rightarrow 20(x^2 + y^2 + 20(x+y) + 20) = 10^2(x+y+2)^2$$

$$\Rightarrow 4x^2 + 4y^2 + 10xy = 0$$

$$\Rightarrow 2x^2 + 2y^2 + 5xy = 0$$

136. (c)  $a = 25, d = 22 - 25 = -3$ .

Let  $n$  be the no. of terms

$$\text{Sum} = 116; \text{Sum} = \frac{n}{2} [2a + (n-1)d]$$

$n$

$$116 = \frac{n}{2} [50 + (n-1)(-3)] \text{ or } 232 = n[50 - 3n + 3] = n[53 - 3n]$$

$$= -3n^2 + 53n$$

$$\Rightarrow 3n^2 - 53n + 232 = 0 \Rightarrow (n-8)(3n-29) = 0$$

$$\Rightarrow n = 8 \text{ or } n = \frac{29}{3}, n \in \mathbb{N} \Rightarrow n = 8$$

$$\backslash \text{Now, } T_8 = a + (8-1)d = 25 + 7 \times (-3) = 25 - 21$$

$$\backslash \text{Last term} = 4$$

137. (d)  $2a = 1 + P$  and  $g^2 = P$

$$P = -P + g^2 \Rightarrow 2P = g^2 \Rightarrow 2a = g^2 \Rightarrow a = \frac{g^2}{2}$$

138. (b)  $y = \sin x = e^x dy^x$

$$P \frac{dx}{dy} = \cos x + e dx$$

$$\frac{dx}{dy} = \frac{1}{\cos x} + e dx \dots (i)$$

$$\backslash \frac{dy}{dx} = -(\cos x + e)^2 [-\sin x + e] \frac{dx}{dy}$$

$$\frac{(e^x - \sin x)x}{(\cos x + e)^3} \cdot \frac{1}{\cos x + e}$$

$$= -\frac{(e^x - \sin x)}{(\cos x + e)^3} \cdot \frac{\sin x e^{-x}}{(\cos x + e)^3}$$

139. (b)  $4x^2 - 9y^2 = 1 \Rightarrow \frac{x^2}{\frac{1}{4}} - \frac{y^2}{\frac{1}{9}} = 1$

$$\frac{x^2}{\frac{1}{4}} - \frac{y^2}{\frac{1}{9}} = 1$$

$$a^2 = \frac{1}{4}, b^2 = \frac{1}{9}$$

$$\frac{a^2}{b^2} = \frac{9}{4} \Rightarrow \frac{3}{2}$$


$$e^2 = \frac{3}{2}$$

$$\text{eccentricity, } e = \sqrt{1 + \frac{b^2}{a^2}} = \sqrt{1 + \frac{1/9}{1/4}} = \sqrt{1 + \frac{4}{9}} = \sqrt{\frac{13}{9}} = \frac{\sqrt{13}}{3}$$

$$= \frac{\sqrt{13}}{3}$$

$$\begin{aligned} \text{æ } 1 \quad 13 \text{ ö } \text{æ } \quad 13 \\ \text{ö foci} = \text{ççè} \pm 2' 3, 0 \div \div \emptyset = \text{ççè} \pm \\ 6, 0 \div \div \emptyset \end{aligned}$$

140. (d) Let height of the tower be  $h$  m and distance between tower and cliff be  $x$  m.  $45^\circ$



$$\setminus CD = h, BD = x$$

$$\text{In } \triangle DAB, \tan 45^\circ = \frac{AB}{BD} \text{ or } 1 = \frac{50}{x}$$

$$x = 50 \quad \dots (i)$$

In  $\triangle AEC$

$$\tan 30^\circ = \frac{AE}{EC} = \frac{AB - EB}{EC} = \frac{AB - DC}{BD}$$

$$(\because EB = DC, EC = BD)$$

$BD$ )

$$\frac{1}{\sqrt{3}} = \frac{50 - h}{x} \text{ or } x = 50 - h \sqrt{3}$$

$$50 = 50 - h \sqrt{3} \quad [\text{From (i), } x = 50]$$

$$\text{or } h = \frac{50(\sqrt{3}-1)}{\sqrt{3}} = 50 \frac{1-\frac{1}{\sqrt{3}}}{\sqrt{3}} = 13.0 \div \div \emptyset$$

$$\setminus h = 50 \frac{1-\frac{1}{\sqrt{3}}}{\sqrt{3}} = 13.0 \div \div \emptyset$$

141. (a) General term of the given binomial series is given by:

$$\begin{aligned} T_{r+1} &= 10C_r \left(\frac{x}{2}\right)^r (10-r)^{-r} \cdot \left(\frac{x}{4}\right)^{1/3} \\ \text{Put } r &= 4, \text{ we get} \end{aligned}$$

$$T_5 = 10C_4 \left(\frac{x}{2}\right)^4 (10-4)^{-4} \cdot \left(\frac{x}{4}\right)^{1/3}$$

$$= \frac{10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{4 \cdot 3 \cdot 2 \cdot 1 \cdot 3} \cdot \frac{70}{243} x^2$$

$$2 = 70 \cdot \frac{\dots}{243}$$

Thus coefficient of  $x$

142. (c) Two circles  $x^2 + y^2 + 2g_1x + 2f_1y + c_1 = 0$  and  $x^2 + y^2 + 2g_2x + 2f_2y + c_2 = 0$  cuts orthogonally if  $2g_1g_2 + 2f_1f_2 = c_1 + c_2$  Given equations of two circles are  $x^2 + y^2 + 2lx + 6y + 1 = 0 \dots (i)$

$$x^2 + y^2 + 4x + 2y = 0 \quad \dots (ii)$$

On comparing (i) and (ii) with original equation, we get

$$g_1 = l, f_1 = 3, c_1 = 1 \text{ and } g_2 = 2, f_2 = 1, c_2 = 0$$

So, from orthogonality condition, we have

$$4l + 6 = 1 + 4 \implies l = -5$$

$$\setminus l = \frac{-5}{4}$$

143. (a) We know, scalar triple product

$$[a \ b \ c] = a \cdot (b \times c) = (a \times b) \cdot c$$

Consider  $[a \ b \ b \ c \ c \ a \ + \ + \ +]$

$$= (a \ b \ +) \cdot \{(b \ c \ +)' (c \ a \ +)\}$$

$$= (a \ + \ b) \cdot \{(b \ c') + (b \ a') + (c \ c') + (c \ a')\}$$

$$= (a \ b \ +) \cdot \{(b \ c') + (b \ a') + (c \ a')\}$$

$$(\because c' \cdot c = 0)$$

$$\begin{aligned}
 & \dots \dots \dots = a b c a b a a \\
 & c a. ( ' + ) . ( ' + ) . ( ' ) \\
 & \dots \dots \dots + b b c b b a b c a. ( ' + ) . ( ' \\
 & + ) . ( ' ) \dots \dots \dots \\
 & = [a b c] + [a b a] + [a c a] + [b b c] \\
 & \dots \dots \dots \\
 & + [b b a] + [b c a] \\
 & \text{(By definition of scalar triple product)} \\
 & \dots \dots \dots \\
 & [a a b] = 0, [a b a] = 0 \text{ and } [b a a] = \\
 & 0 \dots \dots \dots \\
 & = [a b c] + [b c a] = 2 [a b c]
 \end{aligned}$$

144. (b) Given that  $f(x) = (a - x^{n/1/n})$   
 $\setminus \text{fof}(x) = [a - \{(a - x^{n/1/n})^{1/n}\}]$   
 $= [a - (a - x^n)]^{1/n}$   
 $= [x^n]^{1/n} = x$

145. (a) Sum =  $\frac{8}{9} [9 + 99 + 999 + \dots n \text{ terms}]$   
 $= \frac{8}{9} [(10-1) + (100-1) + (1000-1) + \dots n \text{ terms}]$   
 $= \frac{8}{9} [(10 + 10^2 + 10^3 + \dots + 10^n) - n]$   
 $= \frac{8}{9} \left[ \frac{10(10^n - 1)}{10 - 1} - n \right]$   
 $= \frac{8}{81} [10^{n+1} - 9n - 10]$

146. (a) Let  $z = x + iy$   
 $\setminus |z + 3 - i| = |(x + 3) + i(y - 1)| = 1$   
 $\rho \sqrt{(x + 3)^2 + (y - 1)^2} = 1 \dots (i)$   
 $\therefore \arg z = \rho \rho \tan^{-1} y = \rho x$

$$\begin{aligned}
 & y \\
 \rho \quad & \_ = \tan \rho = 0 \quad \rho \quad y = 0 \quad \dots (ii) \\
 & x
 \end{aligned}$$

From equations (i) and (ii), we get  $x = -3, y = 0 \setminus z = -3 \rho |z| = |-3| = 3$

147. (b) Let  $E_1, E_2$  and  $A$  be the events defined as follows:

$E_1$  = red ball is transferred from bag P to bag Q

$E_2$  = blue ball is transferred from bag P to bag Q

$A$  = the ball drawn from bag Q is blue

As the bag P contains 6 red and 4 blue balls,

$$\begin{aligned}
 P E(1) &= \frac{6}{10} \frac{3}{5} \text{ and } P E(2) = \frac{4}{10} \frac{2}{5}
 \end{aligned}$$

Note that  $E_1$  and  $E_2$  are mutually exclusive and exhaustive events.

When  $E_1$  has occurred i.e., a red ball has already been transferred from bag P to Q, then bag Q will contain 6 red and 6 blue

balls, So,  $P(A|E_1) = \frac{6}{12} = \frac{1}{2}$

When  $E_2$  has occurred i.e., a blue ball has already been transferred from bag P to Q, then bag Q will contain 5 red and 7 blue

balls, So,  $P(A|E_2) = \frac{7}{12}$

By using law of total probability, we get


$$P(A) = P(E_1) P(A|E_1) + P(E_2) P(A|E_2)$$

$$\begin{aligned}
 &= \frac{6}{10} \frac{1}{2} + \frac{4}{10} \frac{7}{12} \\
 &= \frac{3}{5} + \frac{7}{15} = \frac{6}{5} + \frac{7}{15} = \frac{12}{15} + \frac{7}{15} = \frac{19}{15}
 \end{aligned}$$

148. (a) Total number of 4-digit numbers  
 $= 5 \times 5 \times 5 \times 5 = 625$   
(as each place can be filled by any one of the numbers 1, 2, 3, 4 and 5)  
Numbers in which no two digits are identical  
 $= 5 \times 4 \times 3 \times 2 = 120$  (i.e. repetition not allowed)

(as 1<sup>st</sup> place can be filled in 5 different ways, 2<sup>nd</sup> place can be filled in 4 different ways and so on)

Number of 4-digits numbers in which at least 2 digits are identical =  $625 - 120 = 505$



149. (c)  $D=2 \quad 3 \quad 1=0 ; D_1=3 \quad 3$   
 $1^1 0$   
 $3 \quad 5 \quad 21 \quad 5 \quad 2$

∅ Given system, does not have any solution. ∅ No solution

150. (d) Let A(3, y), B(2, 7), C(-1, 4) and D(0, 6) be the given points.

$$\text{slope of AB} = \frac{7-y}{2-3} = m_1$$

$$m_2 = \text{slope of CD} = \frac{6-4}{0-(-1)} = 2$$

Since AB and CD are parallel.  $\therefore m_1 = m_2 \Rightarrow y = 9$ .